

### Nutrition Care Indicator Categories

Nutrition Care Indicator Category	Priority Points: <b>4</b>	Priority Points: <b>3</b>	Priority Points: <b>2</b>	Priority Points: <b>0</b>
<b>Nutritional Hx or anticipated</b>	<ul style="list-style-type: none"> <li>▪ No appetite (&lt;50% for &gt;5days)</li> </ul>	<ul style="list-style-type: none"> <li>▪ Poor appetite (50% of meals for &gt; 5days)</li> <li>▪ Diarrhea</li> <li>▪ Swallowing problems</li> <li>▪ Vomiting</li> </ul>	<ul style="list-style-type: none"> <li>▪ Fair appetite (75% of meals for &gt;5 days)</li> <li>▪ Eating, chewing problems</li> <li>▪ Nausea</li> </ul>	<ul style="list-style-type: none"> <li>▪ Good appetite</li> </ul>
<b>Feeding Modality or anticipated</b>	<ul style="list-style-type: none"> <li>▪ TPN/PPN</li> <li>▪ Unstable tube feeding</li> <li>▪ NPO/unsupplemented Clear Liquid Diet &gt;72 hours</li> <li>▪ Unsupplemented Full Liquid Diet &gt;7 days</li> <li>▪ Calorie Count</li> </ul>	<ul style="list-style-type: none"> <li>▪ Diet providing &lt;1000 kcal</li> <li>▪ Restrictive therapeutic diet</li> <li>▪ Multiple allergies</li> <li>▪ Stable tube feeding</li> </ul>	<ul style="list-style-type: none"> <li>▪ Mineral restriction other than sodium</li> </ul>	<ul style="list-style-type: none"> <li>▪ All other diet orders</li> </ul>
<b>Unintentional weight loss/gain</b>	<ul style="list-style-type: none"> <li>▪ &gt;15% in 6 months, or</li> <li>▪ 5% in one month</li> </ul>	<ul style="list-style-type: none"> <li>▪ 10-15% in 6 months or</li> <li>▪ &lt;5% in 1 month</li> </ul>	<ul style="list-style-type: none"> <li>▪ &lt;10% - 6 months</li> </ul>	<ul style="list-style-type: none"> <li>▪ Stable weight</li> </ul>
<b>Weight Status</b>	<ul style="list-style-type: none"> <li>▪ ≤74% IBW</li> </ul>	<ul style="list-style-type: none"> <li>▪ 80-75% IBW</li> </ul>	<ul style="list-style-type: none"> <li>▪ 89-81% IBW</li> </ul>	<ul style="list-style-type: none"> <li>▪ 100-90% IBW</li> </ul>
<b>Dx/Condition</b>	<ul style="list-style-type: none"> <li>▪ Acute renal failure</li> <li>▪ AIDS</li> <li>▪ Burns (&gt;25% BSA)</li> <li>▪ Cachexia</li> <li>▪ Coma or Hepatic Coma</li> <li>▪ GI Obstruction</li> <li>▪ Head or Neck CA</li> <li>▪ Hepatic encephalopathy</li> <li>▪ Hyperemesis Gravidarium</li> <li>▪ Ileus</li> <li>▪ Metastatic CA</li> <li>▪ Multiple trauma</li> <li>▪ Multi-system organ failure</li> <li>▪ Peritonitis</li> <li>▪ Pressure ulcer, stage IV or multiple ulcers</li> <li>▪ Pulmonary failure (vent dependent)</li> <li>▪ Sepsis</li> <li>▪ Transplantation, major organ system, Pre- and Post-op</li> </ul>	<ul style="list-style-type: none"> <li>▪ Burns (&lt;25% BSA)</li> <li>▪ CHF with complications</li> <li>▪ COPD, unstable</li> <li>▪ Chronic renal failure</li> <li>▪ Diabetes, uncontrolled or Gestational</li> <li>▪ Diabetes, new Dx</li> <li>▪ Dysphagia</li> <li>▪ Esophageal, Pancreatic or Stomach CA</li> <li>▪ GI disease with malabsorption of maldigestion</li> <li>▪ Hepatic disease</li> <li>▪ Pressure ulcers stage III</li> <li>▪ Spinal Cord Injury – new</li> <li>▪ Pancreatitis</li> <li>▪ Pregnancy and &lt;18 years of age</li> <li>▪ Pregnancy with other medical conditions</li> </ul>	<ul style="list-style-type: none"> <li>▪ &gt;65 years of age and surgery</li> <li>▪ CABG surgery</li> <li>▪ Colon or Rectal CA</li> <li>▪ Fracture – of long bone, hip, pelvis and &gt;65 years old</li> <li>▪ Fractured jaw</li> <li>▪ Morbid obesity</li> <li>▪ Nutritional anemia</li> <li>▪ Pressure ulcer, stage I or II</li> </ul>	<ul style="list-style-type: none"> <li>▪ Compassion care</li> </ul>
<b>Anthropometrics</b>				

Nutritional Status Classification	Points	Time frame for reassessment/follow-up
Normal (Status 1)	0-5	4-5 days
Mildly Compromised (Status 2)	6-7	4-5 days
Moderately Compromised (Status 3)	8-11	3-4 days
Severely Compromised (Status 4)	≥12	1-3 days