#### **Criterion A: Stressor**

The person was exposed to: death, threatened death, actual or threatened serious injury, or actual or threatened sexual violence, as follows: (one required)

- 1. Direct exposure.
- 2. Witnessing, in person.
- 3. Indirectly, by learning that a close relative or close friend was exposed to trauma. If the event involved actual or threatened death, it must have been violent or accidental.
- 4. Repeated or extreme indirect exposure to aversive details of the event(s), usually in the course of professional duties (e.g., first responders, collecting body parts; professionals repeatedly exposed to details of child abuse). This does not include indirect non-professional exposure through electronic media, television, movies, or pictures.

### **Criterion B: Intrusion Symptoms**

The traumatic event is persistently re-experienced in the following way(s): (one required)

- 1. Recurrent, involuntary, and intrusive memories. Note: Children older than six may express this symptom in repetitive play.
- 2. Traumatic nightmares. Note: Children may have frightening dreams without content related to the trauma(s).
- 3. Dissociative reactions (e.g., flashbacks) which may occur on a continuum from brief episodes to complete loss of consciousness. Note: Children may reenact the event in play.
- 4. Intense or prolonged distress after exposure to traumatic reminders.
- 5. Marked physiologic reactivity after exposure to trauma-related stimuli.

### Criterion C: Avoidance

Persistent effortful avoidance of distressing trauma-related stimuli after the event: (one required)

- 1. Trauma-related thoughts or feelings.
- 2. Trauma-related external reminders (e.g., people, places, conversations, activities, objects, or situations)

## Criterion D: Negative Alterations in Cognitions and Mood

Negative alterations in cognitions and mood that began or worsened after the traumatic event: (two required)

- 1. Inability to recall key features of the traumatic event (usually dissociative amnesia; not due to head injury, alcohol, or drugs).
- 2. Persistent (and often distorted) negative beliefs and expectations about oneself or the world (e.g., "I am bad," "The world is completely dangerous").
- 3. Persistent distorted blame of self or others for causing the traumatic event or for resulting consequences.
- 4. Persistent negative trauma-related emotions (e.g., fear, horror, anger, guilt, or shame).
- 5. Markedly diminished interest in (pre-traumatic) significant activities.
- 6. Feeling alienated from others (e.g., detachment or estrangement).
- 7. Constricted affect: persistent inability to experience positive emotions.

# Criterion E: Alterations in Arousal and Reactivity

Trauma-related alterations in arousal and reactivity that began or worsened after the traumatic event: (two required)

- 1. Irritable or aggressive behavior
- 2. Self-destructive or reckless behavior
- 3. Hypervigilance
- 4. Exaggerated startle response
- 5. Problems in concentration
- 6. Sleep disturbance

## **Criterion F: Duration**

Persistence of symptoms (in Criteria B, C, D, and E) for more than one month.

## Criterion G: Functional Significance

Significant symptom-related distress or functional impairment (e.g., social, occupational).

### **Criterion H: Exclusion**

Disturbance is not due to medication, substance use, or other illness.

**Source:** "DSM-5 Criteria for PTSD." *DSM-5 Criteria for PTSD | BrainLine.* BrainLine, 10 June 2014. Web. <a href="https://www.brainline.org/article/dsm-5-criteria-ptsd">https://www.brainline.org/article/dsm-5-criteria-ptsd</a>.